

References

- Lemerle, D. Gill, G.S. Murphy, C.E. Walker, S.R., Cousens, R.D. Mokhtari, S., Peltzer, S.J., Coleman, R., Luckett, D.J., 2001. Genetic improvement and agronomy for enhanced wheat competitiveness with weeds. *Aust. J. Agric. Res.* 52, 527-548.
- Munier-Jolain, N.M., Chauvel, B., Gasquez, J., 2004. Stratégies de protection intégrée contre les adventices des cultures: le retour de l'agronomie. In: Regnault-Roger, C. (Ed.), *Enjeux Phytosanitaires pour l'Agriculture et l'Environnement du XXI^{ème} Siècle*, Lavoisier, Paris, pp. 411-430.
- Olsen, J., Kristensen, L., Weiner, J., Griepentrog, H.W., 2005. Increased density and spatial uniformity increase weed suppression by spring wheat. *Weed Res.* 45, 361-321.
- Rasmussen, I.A., 2004. The effect of sowing date, stale seedbed, row width and mechanical weed control on weeds and yields of organic winter wheat. *Weed Res.* 44, 12-20.
- REFERENCE: (Chikowo, R., Faloya, V., Petit, S., & Munier-Jolain, N.M. (2009). Integrated weed management systems allow reduced reliance on herbicides and long-term weed control. *Agriculture, Ecosystems and Environment*, 132, 237-242.)